arXiv:1403.3081v1 [math.NT] 12 Mar 2014

EVALUATING BINOMIAL CHARACTER SUMS MODULO POWERS OF TWO

VINCENT PIGNO, CHRIS PINNER, AND JOE SHEPPARD

Abstract. We show that for any mod 2^m characters, χ_1, χ_2, the complete exponential sum,

$$\sum_{x=1}^{2^m} \chi_1(x)\chi_2(Ax^k + B),$$

has a simple explicit evaluation.

1. Introduction

Suppose that χ_1 and χ_2 are mod 2^m multiplicative characters with χ_2 primitive mod 2^m, $m \geq 3$. We are interested here in evaluating the complete character sum

$$S = \sum_{x=1}^{2^m} \chi_1(x)\chi_2(Ax^k + B).$$

Writing $\chi(Ax^K + Bx^L) = \chi^L(x)\chi(Ax^{K-L} + B)$ these sums of course include the binomial character sums. Cases where one can explicitly evaluate an exponential or character sum are unusual and therefore worth investigating.

In [8] we considered the corresponding result for mod p^m characters with $p \geq 3$ and m sufficiently large, using reduction techniques of Cochrane & Zheng [2] (see also [3] & [4]). Though their results are stated for odd primes the approach can often be adapted for $p = 2$ as we showed for twisted monomial exponential sums in [7]. When $k = 1$ and $A = -1$, $B = 1$ the mod p^m sum is the classical Jacobi sum (though uninterestingly zero if $p = 2$). See [1] or [5] for an extensive treatment of mod p Jacobi sums and their generalizations over $\mathbb{F}_{p^m}$. In [6] we treated the $k = 1$ case for general A, B, including when $p = 2$, along with generalizations of the multivariable Jacobi sums considered in [9].

Plainly $S = 0$ if A and B are not of opposite parity (otherwise x or $Ax^k + B$ will be even and the individual terms will all be zero). We assume here that A is even and B is odd and write

$$A = 2^n A_1,\ n > 0, \quad k = 2^t k_1, \quad 2 \nmid A_1 k_1 B.$$

If B is even and A odd we can use $x \mapsto x^{-1}$ to write S in the form

$$S = \sum_{x=1}^{2^m} \overline{\chi_1\chi_2}^k(x)\chi_2(Bx^k + A).$$

Date: March 13, 2014.

2010 *Mathematics Subject Classification.* Primary: 11L05; Secondary: 11L03, 11L10.

Key words and phrases. Character sums.

The first and third authors acknowledge support of K-State's I-Center and Arts & Sciences Undergraduate Research Scholarships respectively.

Since $\mathbb{Z}_{2^m}^* =< -1, 5 >$, the characters χ_1, χ_2 are completely determined by their values on -1 and 5. Since 5 has order 2^{m-2} mod 2^m we can define integers c_1, c_2 with

$$\chi_i(5) = e_{2^{m-2}}(c_i), \quad 1 \le c_i \le 2^{m-2},$$

where $e_n(x) := e^{2\pi i x/n}$. Since χ_2 is primitive we have $2 \nmid c_2$. We define the odd integers R_i, $i \ge 2$, by

$$5^{2^{i-2}} = 1 + R_i 2^i. \tag{1}$$

Defining

$$N := \begin{cases} \lceil \frac{1}{2}(m-n) \rceil, & \text{if } m-n > 2t+4, \\ t+2, & \text{if } t+2 \le m-n \le 2t+4, \end{cases}$$

and

$$C(x) := c_1(Ax^k + B) + c_2 A k x^k R_N R_{N+n}^{-1} \tag{2}$$

(here and throughout the paper y^{-1} denotes the inverse of y mod 2^m) it transpires that the sum S will be zero unless there is a solution x_0 to the characteristic equation

$$C(x_0) \equiv 0 \bmod 2^{\lfloor \frac{1}{2}(m+n) \rfloor + t}, \tag{3}$$

with $2 \nmid x_0(Ax_0^k + B)$, when $m - n > 2t + 4$, and a solution to $C(1)$ or $C(-1) \equiv 0$ mod 2^{m-2} when $t + 2 \le m - n \le 2t + 4$.

Theorem 1.1. *Suppose that $m - n \ge t + 2$. The sum $S = 0$ unless $c_1 = 2^{n+t}c_3$, with $2 \nmid c_3$, and $\chi_1(-1) = 1$ when k is even, and the characteristic equation (3) has an odd solution x_0 when $m - n > 2t + 4$. Assuming these conditions do hold.*

When $m - n > 2t + 4$,

$$S = 2^{\frac{1}{2}(m+n)+t+\min\{1,t\}} \chi_1(x_0)\chi_2(Ax_0^k + B) \begin{cases} 1, & \text{if } m-n \text{ is even}, \\ \omega^h \left(\frac{2}{h}\right), & \text{if } m-n \text{ is odd}, \end{cases}$$

where $\left(\frac{2}{x}\right)$ is the Jacobi symbol, $\omega = e^{\pi i/4}$, $C(x_0) = \lambda 2^{\lfloor \frac{1}{2}(m+n) \rfloor + t}$ for some integer λ and $h := 2\lambda + (k_1 - 1) + (2^n - 1)c_3$.

When $t + 3 < m - n \le 2t + 4$,

$$S = \begin{cases} 2^{m-1}\chi_2(A+B), & \text{if } k \text{ is even and } C(1) \equiv 0 \bmod 2^{m-2}, \\ 2^{m-2}\chi_2(A+B), & \text{if } k \text{ is odd and } C(1) \equiv 0 \bmod 2^{m-2}, \\ 2^{m-2}\chi_1(-1)\chi_2(-A+B), & \text{if } k \text{ is odd and } C(-1) \equiv 0 \bmod 2^{m-2}, \\ 0, & \text{otherwise.} \end{cases}$$

When $m - n = t + 3$,

$$S = \begin{cases} 2^{m-1}\chi_2(A+B), & \text{if } k \text{ is even and } \chi_1(5) = \pm 1,\ \chi_1(-1) = 1, \\ 2^{m-2}\left(\chi_2(A+B) + \chi_1(-1)\chi_2(-A+B)\right), & \text{if } k \text{ is odd and } \chi_1(5) = \pm 1, \\ 0, & \text{otherwise.} \end{cases}$$

When $m - n = t + 2$,

$$S = \begin{cases} 2^{m-1}\chi_2(A+B), & \text{if } k \text{ is even and } \chi_1 = \chi_0 \text{ or } k \text{ is odd and } \chi_1 = \chi_4, \\ 0, & \text{otherwise,} \end{cases}$$

where χ_0 is the principal character mod 2^m and χ_4 is the mod 2^m character induced by the non-trivial character mod 4 (i.e. $\chi_4(x) = \pm 1$ as $x \equiv \pm 1$ mod 4 respectively).

Note that the restriction $m-n \geq t+2$ is quite natural; for $m-n < t+2$ the odd x have $Ax^k+B \equiv A+B \bmod 2^m$ and $S = \chi_2(A+B)\sum_{x=1}^{2^m}\chi_1(x) = 2^{m-1}\chi_2(A+B)$ if $\chi_1 = \chi_0$ and zero otherwise.

Our original assumption that χ_2 is primitive is also reasonable; if χ_1 and χ_2 are both imprimitive then one should reduce the modulus, while if χ_1 is primitive and χ_2 imprimitive then $S = 0$ (if χ_1 is primitive then $u = 1+2^{m-1}$ must have $\chi_1(u) = -1$, since $x + 2^{m-1} \equiv ux \bmod 2^m$ for any odd x, and $x \mapsto xu$ gives $S = \chi_1(u)S$ when χ_2 is imprimitive).

2. Proof

Initial decomposition

Observing that $\pm 5^\gamma$, $\gamma = 1, \dots, 2^{m-2}$, gives a reduced residue system mod 2^m and writing

$$S(A) := \sum_{\gamma=1}^{2^{m-2}} \chi_1(5^\gamma)\chi_2(A5^{\gamma k}+B),$$

if k is even we have

$$S = (1+\chi_1(-1))S(A) = \begin{cases} 0, & \text{if } \chi_1(-1) = -1, \\ 2S(A), & \text{if } \chi_1(-1) = 1, \end{cases} \tag{4}$$

and if k is odd

$$S = S(A) + \chi_1(-1)S(-A). \tag{5}$$

Large m values: $m > n+2t+4$

If I_1 is an interval of length $2^{\lceil \frac{m-n}{2} \rceil - t - 2}$ then plainly

$$\gamma = u2^{\lceil \frac{m-n}{2} \rceil - t - 2} + v, \quad v \in I_1, \quad u \in I_2 := \left[1, 2^{\lfloor \frac{m+n}{2} \rfloor + t}\right],$$

runs through a complete set of residues mod 2^{m-2}. Hence, writing $h(x) := Ax^k+B$ and noting that $2 \nmid h(5^v)$,

$$\begin{aligned} S(A) &= \sum_{v \in I_1} \chi_1(5^v) \sum_{u \in I_2} \chi_1\left(5^{u2^{\lceil \frac{m-n}{2} \rceil - t - 2}}\right) \chi_2\left(A5^{vk}5^{ku2^{\lceil \frac{m-n}{2} \rceil - t - 2}} + B\right) \\ &= \sum_{v \in I_1} \chi_1(5^v)\chi_2(h(5^v)) \sum_{u \in I_2} \chi_1\left(5^{u2^{\lceil \frac{m-n}{2} \rceil - t - 2}}\right) \chi_2(W) \end{aligned}$$

where

$$W = h(5^v)^{-1}A5^{vk}\left(5^{ku2^{\lceil \frac{m-n}{2} \rceil - t - 2}} - 1\right) + 1.$$

Since $n + 2\lceil\frac{m-n}{2}\rceil \geq m$ and $2\lceil\frac{m+n}{2}\rceil \geq m$ we have

$$\begin{aligned}
W &= A_1 5^{vk} h(5^v)^{-1} 2^n \left(\left(1 + R_{\lceil\frac{m-n}{2}\rceil} 2^{\lceil\frac{m-n}{2}\rceil}\right)^{uk_1} - 1\right) + 1 \\
&\equiv 1 + A_1 5^{vk} h(5^v)^{-1} u k_1 R_{\lceil\frac{m-n}{2}\rceil} 2^{\lceil\frac{m+n}{2}\rceil} \bmod 2^m \\
&\equiv \left(1 + R_{\lceil\frac{m+n}{2}\rceil} 2^{\lceil\frac{m+n}{2}\rceil}\right)^{A_1 5^{vk} h(5^v)^{-1} u k_1 R_{\lceil\frac{m-n}{2}\rceil} R^{-1}_{\lceil\frac{m+n}{2}\rceil}} \bmod 2^m \\
&= 5^{A_1 5^{vk} h(5^v)^{-1} u k_1 R_{\lceil\frac{m-n}{2}\rceil} R^{-1}_{\lceil\frac{m+n}{2}\rceil} 2^{\lceil\frac{m+n}{2}\rceil - 2}} \\
&= 5^{A 5^{vk} h(5^v)^{-1} u k R_{\lceil\frac{m-n}{2}\rceil} R^{-1}_{\lceil\frac{m+n}{2}\rceil} 2^{\lceil\frac{m-n}{2}\rceil - t - 2}}.
\end{aligned}$$

So we can write

$$\sum_{u \in I_2} \chi_1\left(5^{u 2^{\lceil\frac{m-n}{2}\rceil - t - 2}}\right) \chi_2(W) = \sum_{u \in I_2} e_{2^{\lfloor\frac{m+n}{2}\rfloor + t}}\left(u\left(c_1 + c_2 A 5^{vk} h(5^v)^{-1} k R_{\lceil\frac{m-n}{2}\rceil} R^{-1}_{\lceil\frac{m+n}{2}\rceil}\right)\right),$$

which equals $2^{\lfloor\frac{m+n}{2}\rfloor + t}$ for the v with

$$c_1 h(5^v) + c_2 A 5^{vk} k R_{\lceil\frac{m-n}{2}\rceil} R^{-1}_{\lceil\frac{m+n}{2}\rceil} \equiv 0 \bmod 2^{\lfloor\frac{m+n}{2}\rfloor + t} \tag{6}$$

and zero otherwise. Since $m \geq n+2$ equation (6) has no solution (and hence $S = 0$) unless $c_1 = 2^{n+t} c_3$ with $2 \nmid c_3$, in which case (6) becomes

$$\left(c_3 A + c_2 A_1 k_1 R_{\lceil\frac{m-n}{2}\rceil} R^{-1}_{\lceil\frac{m+n}{2}\rceil}\right) 5^{vk} \equiv -c_3 B \bmod 2^{\lfloor\frac{m-n}{2}\rfloor}. \tag{7}$$

If no v satisfies (6) then plainly $S = 0$. So assume that (6) has a solution $v = v_0$ and take $I_1 = [v_0, v_0 + 2^{\lceil\frac{m-n}{2}\rceil - t - 2})$. Now any other v solving (7) must have

$$5^{vk} \equiv 5^{v_0 k} \bmod 2^{\lfloor\frac{m-n}{2}\rfloor} \Rightarrow vk \equiv v_0 k \bmod 2^{\lfloor\frac{m-n}{2}\rfloor - 2} \Rightarrow v \equiv v_0 \bmod 2^{\lfloor\frac{m-n}{2}\rfloor - t - 2}.$$

So if $m - n$ is even, I_1 contains only the solution v_0 and

$$S(A) = 2^{\lfloor\frac{m+n}{2}\rfloor + t} \chi_1(5^{v_0}) \chi_2(A 5^{v_0 k} + B). \tag{8}$$

Observe that a solution $x_0 = 5^{v_0}$ or $x_0 = -5^{v_0}$ of (3) corresponds to a solution v_0 to (6) when k is even and a solution v_0 to (6) for A or $-A$ respectively (both can not have solutions) if k is odd. The evaluation for S follows at once from (8) and (4) or (5). When $m - n$ is odd, I_1 contains two solutions v_0 and $v_0 + 2^{\lfloor\frac{m-n}{2}\rfloor - t - 2}$ and

$$\begin{aligned}
S(A) &= 2^{\lfloor\frac{m+n}{2}\rfloor + t} \chi_1(5^{v_0}) \left(\chi_2(h(5^{v_0})) + \chi_1(5^{2^{\lfloor\frac{m-n}{2}\rfloor - t - 2}}) \chi_2(A 5^{v_0 k} 5^{k 2^{\lfloor\frac{m-n}{2}\rfloor - t - 2}} + B)\right) \\
&= 2^{\lfloor\frac{m+n}{2}\rfloor + t} \chi_1(5^{v_0}) \chi_2(h(5^{v_0})) \left(1 + \chi_1(5^{2^{\lfloor\frac{m-n}{2}\rfloor - t - 2}}) \chi_2(\xi)\right)
\end{aligned}$$

where, since $3\lfloor\frac{m-n}{2}\rfloor+n\geq m$ for $m\geq n+3$,

$$\begin{aligned}
\xi &= A5^{v_0k}\left(5^{k_1 2^{\lfloor\frac{m-n}{2}\rfloor-2}}-1\right)h(5^{v_0})^{-1}+1\\
&= A5^{v_0k}h(5^{v_0})^{-1}\left((1+R_{\lfloor\frac{m-n}{2}\rfloor}2^{\lfloor\frac{m-n}{2}\rfloor})^{k_1}-1\right)+1\\
&\equiv A5^{v_0k}h(5^{v_0})^{-1}\left(k_1R_{\lfloor\frac{m-n}{2}\rfloor}2^{\lfloor\frac{m-n}{2}\rfloor}+\binom{k_1}{2}R^2_{\lfloor\frac{m-n}{2}\rfloor}2^{m-n-1}\right)+1 \bmod 2^m\\
&\equiv \left(A_15^{v_0k}h(5^{v_0})^{-1}k_1R_{\lfloor\frac{m-n}{2}\rfloor}R^{-1}_{\lfloor\frac{m+n}{2}\rfloor}+\frac{1}{2}(k_1-1)2^{\lfloor\frac{m-n}{2}\rfloor}\right)R_{\lfloor\frac{m+n}{2}\rfloor}2^{\lfloor\frac{m+n}{2}\rfloor}+1 \bmod 2^m\\
&\equiv \left(1+R_{\lfloor\frac{m+n}{2}\rfloor}2^{\lfloor\frac{m+n}{2}\rfloor}\right)^{A_15^{v_0k}h(5^{v_0})^{-1}k_1R_{\lfloor\frac{m-n}{2}\rfloor}R^{-1}_{\lfloor\frac{m+n}{2}\rfloor}+\frac{1}{2}(k_1-1)2^{\lfloor\frac{m-n}{2}\rfloor}} \bmod 2^m\\
&= 5^{\left(A_15^{v_0k}h(5^{v_0})^{-1}k_1R_{\lfloor\frac{m-n}{2}\rfloor}R^{-1}_{\lfloor\frac{m+n}{2}\rfloor}+\frac{1}{2}(k_1-1)2^{\lfloor\frac{m-n}{2}\rfloor}\right)2^{\lfloor\frac{m+n}{2}\rfloor-2}}.
\end{aligned}$$

Hence, setting

$$c_3+c_2A_15^{v_0k}h(5^{v_0})^{-1}k_1R_{\lceil\frac{m-n}{2}\rceil}R^{-1}_{\lceil\frac{m+n}{2}\rceil}=\lambda 2^{\lfloor\frac{m-n}{2}\rfloor}$$

(only the parity of λ will be used) and recalling that c_2 is odd, we have

$$\begin{aligned}
\chi_1(5^{2^{\lfloor\frac{m-n}{2}\rfloor-t-2}})\chi_2(\xi) &= e_{2^{\lceil\frac{m-n}{2}\rceil}}\left(c_3+c_2A_15^{v_0k}h(5^{v_0})^{-1}k_1R_{\lfloor\frac{m-n}{2}\rfloor}R^{-1}_{\lfloor\frac{m+n}{2}\rfloor}\right)(-1)^{\frac{1}{2}(k_1-1)c_2}\\
&= e_{2^{\lceil\frac{m-n}{2}\rceil}}\left(c_2A_15^{v_0k}h(5^{v_0})^{-1}k_1\left(R_{\lfloor\frac{m-n}{2}\rfloor}R^{-1}_{\lfloor\frac{m+n}{2}\rfloor}-R_{\lceil\frac{m-n}{2}\rceil}R^{-1}_{\lceil\frac{m+n}{2}\rceil}\right)\right)(-1)^{\frac{1}{2}(k_1-1)+\lambda}.
\end{aligned}$$

Since $1+R_{i+1}2^{i+1}=(1+R_i2^i)^2$ we have

$$R_{i+1}=R_i+2^{i-1}R_i^2\equiv R_i+2^{i-1} \bmod 2^{i+2},$$

giving $R_i\equiv 3 \bmod 4$ for $i\geq 3$, and

$$\begin{aligned}
&R_{\lfloor\frac{m-n}{2}\rfloor}R^{-1}_{\lfloor\frac{m+n}{2}\rfloor}-R_{\lceil\frac{m-n}{2}\rceil}R^{-1}_{\lceil\frac{m+n}{2}\rceil}\\
\equiv&R^{-1}_{\lfloor\frac{m+n}{2}\rfloor}R^{-1}_{\lceil\frac{m+n}{2}\rceil}\left((R_{\lceil\frac{m-n}{2}\rceil}-2^{\lceil\frac{m-n}{2}\rceil-2})R_{\lceil\frac{m+n}{2}\rceil}-R_{\lceil\frac{m-n}{2}\rceil}(R_{\lceil\frac{m+n}{2}\rceil}-2^{\lceil\frac{m-n}{2}\rceil+n-2})\right) \bmod 2^{\lceil\frac{m-n}{2}\rceil}\\
\equiv&(1-2^n)2^{\lceil\frac{m-n}{2}\rceil-2} \bmod 2^{\lceil\frac{m-n}{2}\rceil}.
\end{aligned}$$

From (6) we have $c_2A_15^{v_0k}h(5^{v_0})^{-1}k_1\equiv -c_3 \bmod 4$ and

$$S(A)=2^{\lfloor\frac{m+n}{2}\rfloor+t}\chi_1(5^{v_0})\chi_2(h(5^{v_0}))\left(1+i^{(2^n-1)c_3}(-1)^{\frac{1}{2}(k_1-1)+\lambda}\right).$$

The result follows on writing

$$\frac{1+i^h}{\sqrt{2}}=\omega^h\left(\frac{2}{h}\right).$$

Small m values: $t+2\leq m-n\leq 2t+4$

Since $n+2(t+2)\geq m$ we have

$$\begin{aligned}
A5^{\gamma k}+B &= A_12^n(1+R_{t+2}2^{t+2})^{\gamma k_1}+B\\
&\equiv (A+B)\left(1+\gamma k_1A_1R_{t+2}(A+B)^{-1}2^{t+n+2}\right) \bmod 2^m\\
&\equiv (A+B)\left(1+R_{t+n+2}2^{t+n+2}\right)^{\gamma k_1A_1(A+B)^{-1}R_{t+2}R^{-1}_{t+n+2}} \bmod 2^m\\
&= (A+B)5^{\gamma Ak(A+B)^{-1}R_{t+2}R^{-1}_{t+n+2}}.
\end{aligned}$$

Hence $\chi_1(5^\gamma)\chi_2(A5^{\gamma k}+B)$ equals

$$\chi_2(A+B)e_{2^{m-2}}\left(\gamma\left(c_1(A+B)+c_2AkR_{t+2}R_{t+n+2}^{-1}\right)(A+B)^{-1}\right)$$

and $S(A)=2^{m-2}\chi_2(A+B)$ if $C(1)\equiv 0 \bmod 2^{m-2}$ and 0 otherwise. Since $m-n\geq t+2$ the congruence $C(1)\equiv 0 \bmod 2^{m-2}$ implies $c_1=2^{t+n}c_3$ (with c_3 odd if $m-n>t+2$) and becomes

$$c_3(A+B)+c_2A_1k_1R_{t+2}R_{t+n+2}^{-1}\equiv 0 \bmod 2^{m-n-t-2}. \tag{9}$$

For $m-n=t+2$ or $t+3$ this will automatically hold (for both A and $-A$ when k is odd) and $S=2^{m-1}\chi_2(A+B)$ for k even and $\chi_1(-1)=1$, and

$$S=2^{m-2}(\chi_2(A+B)+\chi_1(-1)\chi_2(-A+B))$$

for k odd. Further for k odd and $m-n=2$ we have $-A+B\equiv(1+2^{m-1})(A+B) \bmod 2^m$ with $\chi_2(1+2^{m-1})=-1$ and $S=2^{m-2}\chi_2(A+B)(1-\chi_1(-1))=2^{m-1}\chi_2(A+B)$ if $\chi_1(-1)=-1$ and zero otherwise. Note when $m-n=t+2$ we have $c_1=2^{m-2}$ and $\chi_1(5)=1$ and when $m-n=t+3$ we have $c_1=2^{m-2}$ or 2^{m-3} and $\chi_1(5)=\pm 1$.

Since c_3B is odd (9) can not hold for both A and $-A$ for $m-n>t+3$ and at most one of $S(A)$ or $S(-A)$ is non-zero. When k is odd the congruence condition for $-A$ becomes $C(-1)\equiv 0 \bmod 2^{m-2}$.

References

[1] B.C. Berndt, R.J. Evans & K.S. Williams, *Gauss and Jacobi Sums*, Canadian Math. Soc. series of monographs and advanced texts, vol. 21, Wiley, New York 1998.

[2] T. Cochrane *Exponential sums modulo prime powers*, Acta Arith. 101 (2002), no. 2, 131-149.

[3] T. Cochrane, Zhiyong Zheng, *Pure and mixed exponential sums*, Acta Arith. 91 (1999), no. 3, 249-278.

[4] T. Cochrane, Zhiyong Zheng, *A survey on pure and mixed exponential sums modulo prime powers*, Number theory for the millennium, I (Urbana, IL, 2000), 273-300, A K Peters, Natick,MA, 2002.

[5] R. Lidl & H. Niederreiter, *Finite Fields*, Encyclopedia of Mathematics and its applications 20, 2nd edition, Cambridge University Press, 1997.

[6] M. Long, V. Pigno & C. Pinner, *Evaluating Prime Power Gauss and Jacobi Sums*, submitted Rocky Mountain J. Math. (http://www.math.ksu.edu/∼ pinner/research.html preprint 39.)

[7] V. Pigno & C. Pinner, *Twisted monomial Gauss sums modulo prime powers*, to appear Functiones et Approximatio. (http://www.math.ksu.edu/∼ pinner/research.html preprint 36.)

[8] V. Pigno & C. Pinner, *Binomial Character Sums Modulo Prime Powers*, submitted to Journal de Théorie des Nombres de Bordeaux. (http://www.math.ksu.edu/∼ pinner/research.html preprint 38.)

[9] W. Zhang & Z. Xu, *On the Dirichlet characters of polynomials in several variables*, Acta Arith. 121 (2006), no. 2, 117-124.

Department of Mathematics, Kansas State University, Manhattan, KS 66506

E-mail address: pignov@math.ksu.edu, pinner@math.ksu.edu & jnsheppa@ksu.edu